

Tylenol (acetaminophen) Dosage Chart

Give the weight appropriate dosage listed below every 4 hours as needed for a temperature > 101 °F.

*****Do not give Tylenol to children under 3 months of age without first consulting a doctor.**

| Weight (pounds) | Infant Drops 80mg / 0.8ml | Childrens Liquid 160mg / 5ml | Chewable Tablets 80mg tablets | Junior Strength 160 mg tablets | For Staff Use: Weight (Kg) |
|-----------------|---|---------------------------------|----------------------------------|-----------------------------------|-------------------------------|
| 6 -7 | 40 mg (0.4 ml)* | 1.4 ml | | | 3 |
| 8-9 | 60 mg (0.6 ml)* | 1.9 ml | | | 4 |
| 10-11 | 75 mg (0.8 ml)* | 2.3 ml | | | 5 |
| 12-13 | 90 mg (0.9 ml)* | 2.8 ml | | | 6 |
| 14-16 | 100 mg (1.0 ml)* | 3.1 ml | | | 7 |
| 17-18 | 120 mg (1.2 ml)* | 3.75 ml | | | 8 |
| 19-22 | 135 mg (1.4 ml)* | 4.2 ml | | | 9 |
| 22-26 | 160 mg (1.6 ml)* | 5 ml (1 tsp) | 2 tabs | 1 tab/cap | 10-12 |
| 28-31 | *DISCONTINUED Limited Availability | 6 ml | 2 tabs | 1 tab/cap | 13-14 |
| 32-35 | | 7 ml (1 ½ tsp) | 3 tabs | 1 ½ tabs/caps | 15-16 |
| 36-39 | | 8 ml | 3 tabs | 1 ½ tabs/caps | 17-18 |
| 40-44 | | 9 ml | 3 tabs | 1 ½ tabs/caps | 19-20 |
| 45-48 | | 10 ml (2 tsp) | 4 tabs | 2 tabs/caps | 21-22 |
| 49-53 | | 11 ml | 4 tabs | 2 tabs/caps | 23-24 |
| 54-57 | | 12 ml (2 ½ tsp) | 5 tabs | 2 ½ tabs/caps | 25-26 |
| 58-62 | | 13 ml | 5 tabs | 2 ½ tabs/caps | 27-28 |
| 63-66 | | 14 ml | 5 tabs | 2 ½ tabs/caps | 29-30 |
| 67-71 | | 15 ml (3 tsp) | 6 tabs | 3 tabs/caps | 31-32 |
| 72-75 | | 16 ml | 6 tabs | 3 tabs/caps | 33-34 |
| 76-80 | | 17 ml (3 ½ tsp) | 7 tabs | 3 ½ tabs/caps | 35-36 |
| 81-84 | | 18 ml | 7 tabs | 3 ½ tabs/caps | 37-38 |
| 85-88 | | 19 ml | 7 tabs | 3 ½ tabs/caps | 39-40 |
| 89-93 | | 20 ml (4 tsp) | 8 tabs | 4 tabs/caps | 41-42 |
| 94-99 | | 20 ml (4 tsp) | 8 tabs | 4 tabs/caps | 43-44 |
| ≥ 100 | 20 ml (4 tsp) | 8 tabs | 4 tabs/caps | ≥ 45 | |

When used together, tylenol and motrin may be alternated every 3 hours.

****Consult a physician before using tylenol if your child has any liver problems.***

Ibuprofen (Motrin, Advil) Dosage Chart

Give the weight appropriate dosage listed below every 6 hours as needed for temperature > 101 °F or for pain.

*****Do not give Ibuprofen if child is less than 6 months old*****

| Weight (pounds) | Infant Drops 50mg / 1.25 ml | Childrens Liquid 100mg / 5ml | For Staff Use Weight (Kg) |
|-----------------|---|---------------------------------|------------------------------|
| 8-9 | 40 mg (1.0 ml)* | 40 mg (2.0 ml) | 4 |
| 10-11 | 50 mg (1.25 ml)* | 50 mg (2.5 ml) | 5 |
| 12-13 | 60 mg (1.5 ml)* | 60 mg (3.0 ml) | 6 |
| 14-16 | 70 mg (1.75ml)* | 70 mg (3.5 ml) | 7 |
| 17-18 | 80 mg (2.0 ml)* | 80 mg (4.0 ml) | 8 |
| 19-22 | 90 mg (2.25ml)* | 90 mg (4.5 ml) | 9 |
| 22-26 | 100 mg (2.5ml)* | 5 ml (1 tsp) | 10-11 |
| 28-31 | 120 mg (3.0ml)* | 6 ml | 12-13 |
| 32-35 | 140 mg (3.5ml)* | 7 ml (1 ½ tsp) | 14-15 |
| 36-39 | 160 mg (4.0 ml)* | 8 ml | 16-17 |
| 40-44 | *DISCONTINUED Limited Availability | 9 ml | 18-19 |
| 45-48 | | 10 ml (2 tsp) | 20-21 |
| 49-53 | | 11 ml | 22-23 |
| 54-57 | | 12 ml (2 ½ tsp) | 24-25 |
| 58-62 | | 13 ml | 26-27 |
| 63-66 | | 14 ml | 28-29 |
| 67-71 | | 15 ml (3 tsp) | 30-31 |
| 72-75 | | 16 ml | 32-33 |
| 76-80 | | 17 ml (3 ½ tsp) | 34-35 |
| 81-84 | | 18 ml | 36-37 |
| 85-88 | | 19 ml | 38-39 |
| 89-93 | | 20 ml (4 tsp) | 40-41 |
| 94-99 | | 20 ml (4 tsp) | 42-43 |
| ≥ 100 | 20 ml (4 tsp) | ≥ 44 | |

When used together, tylenol and motrin may be alternated every 3 hours.

****Consult a physician before using motrin if your child has any kidney problems.***